arXiv:1703.04968v1 [cs.IT] 15 Mar 2017

Several Classes of Trace Codes With Either Optimal Two Weights or a Few Weights over $\mathbb{F}_q + u\mathbb{F}_q$ *

Hongwei Liu[1], Youcef Maouche[1,2]

[1]School of Mathematics and Statistics, Central China Normal University,Wuhan, Hubei, 430079, China
[2]Department of Mathematics, University of Sciences and Technology HOUARI BOUMEDIENE, Alger, Algeria

Abstract. Let p be a prime number, $q = p^s$ for a positive integer s. For any positive divisor e of $q-1$, we construct an infinite families codes of dimension $2m$ with few Lee-weight. These codes are defined as trace codes over the ring $R = \mathbb{F}_q + u\mathbb{F}_q$, $u^2 = 0$. Using Gauss sums, their Lee weight distribution is provided. When $\gcd(e, m) = 1$, we obtain an infinite family of two-weight codes which meet the Griesmer bound. Moreover, if $\gcd(e, m) = 2, 3$ or 4 we construct new infinite families with at most five-weight. These codes can be used in authentication codes and secret sharing schemes.

Keywords. Two-weight codes; Three-weight codes; Codes over rings; Trace codes; Gauss sums.

2010 *Mathematics Subject Classification.* 94B05, 94B15.

1 Introduction

Throughout this paper, let p be a prime, $q = p^s$ for a positive integer s, and $r = q^m$ for a positive integer m. We call C an $[n, k, d]$ linear code over $\mathbb{F}_q$ if it is a k-dimensional subspace of $\mathbb{F}_q^n$ and has minimum Hamming distance d. Let A_i denote the number of codewords with Hamming weight i in a code C of length n. The weight enumerator of a linear code C is defined by $1 + A_1 z + A_2 z^2 + ... + A_n z^n$. The weight distribution $(1, A_1, ..., A_n)$ is an important research topic in coding theory, as it contains crucial information as to estimate the error correcting capability and the probability of error detection and correction with respect to some algorithms. A code C is said to be a t-weight code if the number of nonzero A_i in the sequence $(A_1, A_2, ..., A_n)$ is equal to t. Codes with few weights have been studied in many articles, see for instance [1, 3, 15].

*E-Mail addresses: hwliu@mail.ccnu.edu.cn (H. Liu), m.youcef@mails.ccnu.edu.cn (Y. Maouche).

Let $Tr_{r/q}(\cdot)$ denote the trace function from $\mathbb{F}_r$ to $\mathbb{F}_q$. For a set $D = \{d_1, d_2, ..., d_n\} \subseteq \mathbb{F}_r^*$, define a linear code of length n over $\mathbb{F}_q$ by

$$C_D = \left\{ (Tr_{r/q}(xd_1), Tr_{r/q}(xd_2), ..., Tr_{r/q}(xd_n)) : x \in \mathbb{F}_r \right\}.$$

The set D is called the defining set of C_D. Many well-known linear codes could be produced by selecting the defining set (see for instance [1, 4, 2, 7]). In a series of papers [13, 12, 14], the notion of trace codes has been extended from finite fields to finite rings as follows. Let R be a finite ring, and R_m an extension of R of degree m, a trace code with a defining set $L = \{d_1, d_2, ..., d_n\} \subseteq R_m^*$ is defined by the following formula

$$C_L = \{ (Tr(xd_1), Tr(xd_2), ..., Tr(xd_n)) : x \in R_m \}, \tag{1}$$

where R_m^* is the group of units of R_m, and $Tr(\cdot)$ is a linear function from R_m to R.

The class of finite rings of the form $R = \mathbb{F}_q + u\mathbb{F}_q$ has been used widely as alphabets in certain codes. Throughout this paper we let $R = \mathbb{F}_q + u\mathbb{F}_q$ and $\mathcal{R} = \mathbb{F}_r + u\mathbb{F}_r$ be an extension of R. The Lee weight distribution of the trace code C_L define by (1) is settled for a few special cases and is in general quite complex. The following special cases have been studied in the literature.

- When $L = \mathbb{F}_{2^m}^* + u\mathbb{F}_{2^m}$ and $R = \mathbb{F}_2 + u\mathbb{F}_2$, then the Lee weight distribution of the code C_L is a two-weight code [13].
- When $L = \mathcal{Q} + u\mathbb{F}_{p^m}$ and $R = \mathbb{F}_p + u\mathbb{F}_p$, where p an odd prime and $\mathcal{Q}$ is the set of all square elements of $\mathbb{F}_{p^m}^*$, then C_L is a two-weight or three-weight code [14].

The purpose of this paper is to investigate the case $L = C_0^{(e,r)} + u\mathbb{F}_r$ and $R = \mathbb{F}_q + u\mathbb{F}_q$, where e is a positive divisor of $q-1$ and $C_0^{(e,r)}$ is the cyclotomic class of order e. We also show that our results include the results of [13] as a special case for $q = 2$, our results also include the results of [14] as special cases for $\gcd(e, m) = 2$ and $\gcd(e, m) = 1$. Furthermore, when taking $\gcd(e, m) = 3$ or 4 we construct a new infinite families of codes with at most five-weight.

This paper is organized as follows. The next section sets up some basic notations and definitions, for any positive divisor e of $q-1$, we construct the code $C_q(m, e)$ defined in (4). Section 3 shows that the Lee-weight of the codes $C_q(m, e)$ is related to the value of $\gcd(e, m)$. When $\gcd(e, m) = 1$ the codes defined and investigated in [13] is just a special case for $q = 2$ (see Corollary 3.4), moreover, if $\gcd(e, m) = 1$, our codes achieved the Griesmer bound, and they are optimal for given length and dimension in few cases. We also investigate the cases where $\gcd(e, m) = 2$ and $\gcd(e, m) = 1$ and show that this cases are a generalization of the codes investigated in [14]. Furthermore, if $\gcd(e, m) = 3$ or $\gcd(e, m) = 4$, we construct new infinite families of codes with at most five-weight. These codes can be used in secret sharing schemes (see [13, 14]).

2 Preliminaries

2.1 The Ring $\mathcal{R} = \mathbb{F}_{q^m} + u\mathbb{F}_{q^m}$

Let p be a prime number. The ring $\mathbb{F}_p + u\mathbb{F}_p$ consists of all polynomials of degree 0 and 1 modulo p in indeterminate u, it is closed under addition and multiplication modulo u^2. Thus, $\mathbb{F}_p + u\mathbb{F}_p$ is a local ring with maximal ideal $\langle u \rangle$. For Given integers s and m, we can construct the ring $R = \mathbb{F}_q + u\mathbb{F}_q$ and the extension ring $\mathcal{R} = \mathbb{F}_r + u\mathbb{F}_r$, where $q = p^s$ and $r = q^m$. There is a Frobenius operator F form $\mathcal{R}$ to R which maps $a + ub$ to $a^q + ub^q$. The trace function, denoted by Tr is then defined as

$$Tr = \sum_{j=0}^{m-1} F^j.$$

It is easy to check that

$$Tr(a + ub) = Tr_{r/q}(a) + uTr_{r/q}(b),$$

for all $a, b \in \mathbb{F}_r$. Here $Tr_{r/q}(\cdot)$ denotes the trace function from $\mathbb{F}_r$ to $\mathbb{F}_q$. The ring $\mathcal{R}$ is local with maximal ideal $M = \langle u \rangle$. The residue field $\mathcal{R}/M$ is isomorphic to $\mathbb{F}_r$. The group of units $\mathcal{R}^*$, is isomorphic as a multiplicative group, to the product of a cyclic group of order $r - 1$ by an elementary abelian group of order r.

Let $Tr_{q/p}(\cdot)$ denote the trace function from $\mathbb{F}_q$ to $\mathbb{F}_p$. Let $i = \sqrt{-1}$ be the imaginary unit, an additive character of $\mathbb{F}_q$ is a nonzero function χ from $\mathbb{F}_q$ to the set of complex numbers such that $\chi(x + y) = \chi(x)\chi(y)$ for any pair $x, y \in \mathbb{F}_q$. For each $b \in \mathbb{F}_q$, the function

$$\chi_b(c) = e^{2i\pi Tr_{q/p}(bc)/p}, \qquad \text{for all } c \in \mathbb{F}_q,$$

defines an additive character of $\mathbb{F}_q$. When $b = 0$, $\chi_0(c) = 1$ for all $c \in \mathbb{F}_q$, and it is called the trivial additive character of $\mathbb{F}_q$. The character $\chi_1(c)$ is called the canonical additive character of $\mathbb{F}_q$.

A multiplicative character of $\mathbb{F}_q$ is a nonzero function ψ from $\mathbb{F}_q^*$ to the set of complex numbers such that $\psi(xy) = \psi(x)\psi(y)$ for all pairs $x, y \in \mathbb{F}_q^*$. Let g be a fixed primitive element of $\mathbb{F}_q^*$. For each $j = 0, 1, ..., q - 2$, the function with

$$\psi_j(g^k) = e^{2i\pi jk/(q-1)} \quad \text{for } k = 0, 1, ..., q - 1, \tag{2}$$

defines a multiplicative character of $\mathbb{F}_q$. When $j = 0$, $\psi_0(x) = 1$ for all $x \in \mathbb{F}_q^*$, and is called the trivial multiplicative character of $\mathbb{F}_q$.

Let q be odd and $j = (q - 1)/2$ in (2), then we get a multiplicative character such that

$$\eta(g^k) = \psi_{\frac{q-1}{2}}(g^k) = \begin{cases} 1, & \text{if } k \text{ is even,} \\ -1, & \text{otherwise.} \end{cases}$$

This η is called the quadratic character of $\mathbb{F}_q$. Let ψ be a multiplicative and χ an additive character of $\mathbb{F}_q$ respectively. The Gaussian sum $G(\psi, \chi)$ is defined by

$$G(\psi, \chi) = \sum_{c \in \mathbb{F}_q^*} \psi(c)\chi(c).$$

If $\psi \neq \psi_0$ and $\chi \neq \chi_0$, then the absolute value of $G(\psi, \chi)$ is $q^{1/2}$ ([9]). If $q = p^s$, where p is an odd prime and s is a positive integer, then

$$G(\eta, \chi_1) = \begin{cases} (-1)^{s-1} q^{1/2}, & \text{if } p \equiv 1 \pmod 4, \\ (-1)^{s-1} i^s q^{1/2}, & \text{if } p \equiv 3 \pmod 4. \end{cases} \tag{3}$$

The following result [9] is useful in the sequel.

Lemma 2.1. Let χ be a nontrivial additive character of $\mathbb{F}_q$ with q being odd, and let $f(x) = a_2 x^2 + a_1 x + a_0 \in \mathbb{F}_q[x]$ with $a_2 \neq 0$. Then

$$\sum_{c \in \mathbb{F}_q} \chi(f(c)) = \chi(a_0 - a_1^2 (4a_2)^{-1}) \eta(a_2) G(\eta, \chi).$$

2.2 Cyclotomy and Gaussian periods

Let $r - 1 = nN$ for two positive integers $n > 1$ and $N > 1$, and let α be a fixed primitive element of $\mathbb{F}_r$. Define $C_i^{(N,r)} = \alpha^i \langle \alpha^N \rangle$ for $i = 0, 1, ..., N-1$, where $C_0^{(N,r)} = \langle \alpha^N \rangle$ denotes the subgroup of $\mathbb{F}_r^*$ generated by α^N. The cosets $C_i^{(N,r)}$ are called the cyclotomic classes of order N in $\mathbb{F}_r$.

Lemma 2.2 ([10])**.** Let e be a positive divisor of $q - 1$. Then we have the following multiset equality:

$$\left\{ xy : y \in C_0^{(e,r)}, x \in \mathbb{F}_q^* \right\} = \frac{q-1}{e} \gcd(m, e) * C_0^{(\gcd(m,e),r)},$$

where $\frac{q-1}{e} \gcd(m, e) * C_0^{(\gcd(m,e),r)}$ denotes the multiset in which each element in the set $C_0^{(\gcd(m,e),r)}$ appears in the multiset with multiplicity $\frac{q-1}{e} \gcd(m, e)$.

The Gaussian periods are defined by

$$\eta_i^{(N,r)} = \sum_{x \in C_i^{(N,r)}} \chi(x), \quad i = 0, 1, ..., N-1,$$

where χ is the canonical additive character of $\mathbb{F}_r$.

The values of the Gaussian periods are in general very hard to compute. However, they can be computed in a few cases.

The period polynomials $\psi_{(N,r)}(X)$ are defined by

$$\psi_{(N,r)}(X) = \prod_{i=0}^{N-1} \left(X - \eta_i^{(N,r)} \right).$$

It is known that $\psi^{(N,r)}(X)$ is a polynomial with integer coefficients [11]. We will need the following four lemmas whose proofs can be found in [11].

Lemma 2.3. Let $N = 3$. Let c and d be defined by $4r = c^2 + 27d^2$, $c \equiv 1 \pmod 3$, and, if $p \equiv 1 \pmod 3$, then $\gcd(c, p) = 1$. These restrictions determine c uniquely, and d up to sign. Then we have

$$\psi_{(3,r)}(X) = X^3 + X^2 - \frac{r-1}{3} X - \frac{(c+3)r - 1}{27}.$$

Lemma 2.4. Let $N = 3$. We have the following results on the factorization of $\psi_{(3,r)}(X)$.

(a) If $p \equiv 2 \pmod 3$, then sm is even, and

$$\psi_{(3,r)}(X) = \begin{cases} 3^{-3}(3X+1+2\sqrt{r})(3X+1-\sqrt{r})^2, & \text{if } sm/2 \text{ even}, \\ 3^{-3}(3X+1-2\sqrt{r})(3X+1+\sqrt{r})^2, & \text{if } sm/2 \text{ odd}. \end{cases}$$

(b) If $p \equiv 1 \pmod 3$, and $sm \not\equiv 0 \pmod 3$, then $\psi_{(3,r)}(X)$ is irreducible over the rationals.

(c) If $p \equiv 1 \pmod 3$, and $sm \equiv 0 \pmod 3$, then

$$\psi_{(3,r)}(X) = \frac{1}{27}(3X+1-c_1 r^{1/3})(3X+1+\frac{1}{2}(c_1+9d_1)r^{1/3}) \cdot (3X+1+\frac{1}{2}(c_1-9d_1)r^{1/3}),$$

where c_1 and d_1 are given by $4p^{m/3} = c_1^2 + 27d_1^2, c_1 \equiv 1 \pmod 3$ and $\gcd(c_1, p) = 1$.

Lemma 2.5. Let $N = 4$. Let u and v be defined by $r = u^2 + 4v^2$, $u \equiv 1 \pmod 4$, and, if $p \equiv 1 \pmod 4$, then $\gcd(u, p) = 1$. These restrictions determine u uniquely, and v up to sign.

If n is even, then

$$\psi_{(4,r)}(X) = X^4 + X^3 - \frac{3r-3}{8}X^2 + \frac{(2u-3)r+1}{16}X + \frac{r^2-(4u^2-8u+6)r+1}{256}.$$

If n is odd, then

$$\psi_{(4,r)}(X) = X^4 + X^3 + \frac{r+3}{8}X^2 + \frac{(2u+1)r+1}{16}X + \frac{9r^2-(4u^2-8u-2)r+1}{256}.$$

Lemma 2.6. Let $N = 4$. We have the following results on the factorization of $\psi_{(4,r)}(X)$.

(a) If $p \equiv 3 \pmod 4$, then sm is even, and

$$\psi_{(4,r)}(X) = \begin{cases} 4^{-4}(4X+1+3\sqrt{r})(4X+1-\sqrt{r})^3, & \text{if } sm/2 \text{ even}, \\ 4^{-4}(4X+1-3\sqrt{r})(4X+1+\sqrt{r})^3, & \text{if } sm/2 \text{ odd}. \end{cases}$$

(b) If $p \equiv 1 \pmod 4$, and sm is odd, then $\psi_{(4,r)}(X)$ is irreducible over the rationals.

(c) If $p \equiv 1 \pmod 4$, and $sm \equiv 2 \pmod 4$, then

$$\psi_{(4,r)}(X) = 4^{-4}((4X+1)^2 + 2\sqrt{r}(4X+1) - r - 2u\sqrt{r})((4X+1)^2 - 2\sqrt{r}(4X+1) - r + 2u\sqrt{r}),$$

the quadratics being irreducible, the u is defined in Lemma 2.5.

(d) If $p \equiv 1 \pmod 4$, and $sm \equiv 0 \pmod 4$, then

$$\begin{aligned}\psi_{(4,r)}(X) = 4^{-4}&((4X+1)+\sqrt{r}+2r^{1/4}u_1)((4X+1)+\sqrt{r}-2r^{1/4}u_1) \\ &\times ((4X+1)-\sqrt{r}+4r^{1/4}v_1)((4X+1)-\sqrt{r}-4r^{1/4}v_1),\end{aligned}$$

where u_1 and v_1 are given by $p^{m/2} = u_1^2 + 4v_1^2, u_1 \equiv 1 \pmod 4$ and $\gcd(u_1, p) = 1$.

The following lemma is well-known, and also can be obtained from Lemma 2.1 and Equation (3).

Lemma 2.7. When $N = 2$, the Gaussian periods are given by the following

$$\eta_0^{(2,r)} = \begin{cases} \frac{-1+(-1)^{sm-1}r^{1/2}}{2}, & \text{if } p \equiv 1 \pmod 4, \\ \frac{-1+(-1)^{sm-1}i^{sm}r^{1/2}}{2}, & \text{if } p \equiv 3 \pmod 4, \end{cases}$$

and $\eta_1^{(2,r)} = -1 - \eta_0^{(2,r)}$.

2.3 Codes and Gray map

A linear code C of length n over $R = \mathbb{F}_q + u\mathbb{F}_q$ is an R-submodule of R^n. If $x = (x_1, x_2, ..., x_n)$ and $y = (y_1, y_2, ..., y_n)$ are two vectors of R^n, the inner product of x and y is defined by $\langle x, y\rangle = \sum_{i=1}^n x_i y_i$, where the operation is performed in R. The dual code of C is denoted by $C^\perp$ and defined as $C^\perp = \{y \in R^n | \langle x, y\rangle = 0, \forall x \in C\}$. $C^\perp$ is also a linear code over R.

The Gray map ϕ from R to $\mathbb{F}_q^2$ is defined by

$$\phi(a + ub) = (b, a + b),$$

for all $a, b \in \mathbb{F}_q$. It is a one to one map from R to $\mathbb{F}_q^2$, which extends naturally into a map from R^n to $\mathbb{F}_q^{2n}$. The Lee weight of a word over R is defined as the Hamming weight of its Gray image

$$w_L(a + ub) = w_H(b) + w_H(a + b)$$

for $a, b \in \mathbb{F}_q^n$. The Lee distance of $x, y \in R^n$ is defined as $w_L(x - y)$. Thus the Gray map by construction is a linear isometry from (R^n, d_L) to $(\mathbb{F}_q^{2n}, d_H)$.

Let e be a positive divisor of $q - 1$ we define $L = C_0^{(e,r)} + u\mathbb{F}_r$. Thus L is a subgroup of $\mathcal{R}^*$ of index e. For $a \in \mathcal{R}$, we define the evaluation map $ev(a)$ by $ev(a) = (Tr(ax))_{x \in L}$ and $C_q(m, e)$ by:

$$C_q(m, e) = \{ev(a) | a \in \mathcal{R}\}. \tag{4}$$

Thus $C_q(m, e)$ is a code over R of length $n = |L| = \frac{r^2 - r}{e}$.

3 The Lee weight distribution

Let χ_1 and χ be the canonical characters of $\mathbb{F}_q$ and $\mathbb{F}_r$ respectively. For any $y \in \mathbb{F}_q$, the following is a basic property of additive characters

$$\sum_{x \in \mathbb{F}_q} \chi_1(xy) = \begin{cases} q, & \text{if } y = 0, \\ 0, & \text{if } y \neq 0. \end{cases} \tag{5}$$

Let N be a positive integer. For $y = (y_1, y_2, ..., y_N) \in \mathbb{F}_q^N$, let

$$\Theta(y) = \sum_{j=1}^{N} \chi_1(y_j).$$

For simplicity, we let $\theta(a) = \Theta(\phi(ev(a)))$, where $a \in \mathcal{R}$. By linearity of the Gray map, and of the evaluation map, we see that $\theta(sa) = \Theta(\phi(ev(sa)))$, for any $s \in \mathbb{F}_q$.

Proposition 3.1. Let N be a positive integer. For all $y = (y_1, y_2, ..., y_N) \in \mathbb{F}_q^N$, we have

$$\sum_{s \in \mathbb{F}_q^*} \Theta(sy) = (q-1)N - qw_H(y).$$

Proof. Using the definition of Θ and Equation (5), we get

$$\sum_{s \in \mathbb{F}_q} \Theta(sy) = \sum_{s \in \mathbb{F}_q} \sum_{j=1}^{N} \chi_1(sy_j) = q \sum_{j=1}^{N} \frac{1}{q} \sum_{s \in \mathbb{F}_q} \chi_1(sy_j) = qN - qw_H(y).$$

Note that

$$\Theta(0) = \sum_{j=1}^{N} \chi_1(0) = N.$$

Hence, the proof follows. □

Proposition 3.2. Let $a = u\beta \in M \setminus \{0\}$ where $M = \langle u \rangle$, then we have

$$\sum_{s \in \mathbb{F}_q^*} \theta(sa) = 2r \frac{q-1}{e} \gcd(e, m) \sum_{s \in C_0^{((e,m),r)}} \chi(\beta s).$$

Proof. We have that

$$\sum_{s \in \mathbb{F}_q^*} \theta(sa) = \sum_{s \in \mathbb{F}_q^*} \Theta\left(\phi\left(ev(as)\right)\right) = \sum_{s \in \mathbb{F}_q^*} \Theta\left(\phi\left(Tr(asx)\right)_{x \in L}\right).$$

Because $a = u\beta$, $x = t + ut'$ with $t' \in \mathbb{F}_q$ and $t \in C_0^{(e,r)}$, then $ax = u\beta t$ and

$$Tr(ax) = Tr(u\beta t) = uTr_{r/q}(\beta t).$$

Let $s \in \mathbb{F}_q^*$, taking Gray map yields

$$\phi(ev(as)) = \phi((Tr(asx))_{x \in L}) = (Tr_{r/q}(\beta st), Tr_{r/q}(\beta st))_{t,t'}.$$

Taking character sum yields

$$\begin{aligned}
\sum_{s \in \mathbb{F}_q^*} \theta(sa) &= \sum_{s \in \mathbb{F}_q^*} \Theta(\phi(ev(as))) = \sum_{s \in \mathbb{F}_q^*} \Theta((Tr_{r/q}(\beta st), Tr_{r/q}(\beta st))_{t,t'}) \\
&= 2 \sum_{s \in \mathbb{F}_q^*} \sum_{t \in C_0^{(r,e)}} \sum_{t' \in \mathbb{F}_r} \chi_1(Tr_{r/q}(\beta st)) \\
&= 2r \sum_{s \in \mathbb{F}_q^*} \sum_{t \in C_0^{(e,r)}} \chi(\beta st).
\end{aligned}$$

Using Lemma 2.2 we get

$$\sum_{s\in\mathbb{F}_q^*}\theta(sa)=2r\frac{q-1}{e}gcd(m,e)\sum_{t\in C_0^{((e,m),r)}}\chi(\beta t). \qquad \square$$

3.1 The Lee weight distribution when $\gcd(e,m)=1$

Theorem 3.3. Let $a\in R$ and $\gcd(e,m)=1$, then the Lee weight of codewords of $C_q(m,e)$ is completely determined by

- If $a=0$, then $w_L(ev(a))=0$,
- If $a\in M\setminus\{0\}$, then $w_L(ev(a))=2\frac{q-1}{eq}r^2$,
- If $a\in\mathcal{R}^*$, then $w_L(ev(a))=2\frac{q-1}{eq}(r^2-r)$.

Proof. If $a=0$, then $Tr(ax)=0$. So $w_L(ev(a))=0$.

If $a=u\beta$ with $\beta\in\mathbb{F}_r^*$, using Proposition 3.2 we get

$$\begin{aligned}\sum_{s\in\mathbb{F}_q^*}\theta(sa)&=2r\frac{q-1}{e}\sum_{s\in C_0^{(1,r)}}\chi(\beta s)\\&=2r\frac{q-1}{e}\sum_{s\in\mathbb{F}_r^*}\chi(\beta s)\\&=-2r\frac{q-1}{e}.\end{aligned}$$

We have that $ev(a)\in R^n$, then $\phi(ev(a))\in\mathbb{F}_q^{2n}$. By Proposition 3.1 we get

$$-2r\frac{q-1}{e}=2(q-1)N-qw_H(\phi(ev(a))),$$

which implies that

$$w_L(ev(a))=2\frac{q-1}{eq}r^2.$$

If $a\in\mathcal{R}^*$ and $x=t+ut'\in L$, then a can be expressed as $a=\alpha+u\beta$. So $ax=\alpha t+u(\alpha t'+\beta t)$, and

$$Tr(ax)=Tr_{r/q}(\alpha t)+uTr_{r/q}(\alpha t'+\beta t),$$

which implies that

$$\phi(ev(a))=\phi((Tr(ax))_{x\in L})=(Tr_{r/q}(\alpha t'+\beta t),Tr_{r/q}(\alpha t'+\beta t)+Tr_{r/q}(\alpha t))_{t,t'}.$$

Taking character sum yields

$$\begin{aligned}\theta(a)&=\sum_{t\in C_0^{(e,r)}}\chi_1(Tr_{r/q}(\beta t))\sum_{t'\in\mathbb{F}_r}\chi_1(Tr_{r/q}(\alpha t'))+\sum_{t\in C_0^{(e,r)}}\chi_1(Tr_{r/q}((\alpha+\beta)t))\sum_{t'\in\mathbb{F}_r}\chi_1(Tr_{r/q}(\alpha t'))\\&=\sum_{t\in C_0^{(e,r)}}\chi(\beta t)\sum_{t'\in\mathbb{F}_r}\chi(\alpha t')+\sum_{t\in C_0^{(e,r)}}\chi((\alpha+\beta)t)\sum_{t'\in\mathbb{F}_r}\chi(\alpha t').\end{aligned}$$

Since $\sum_{t' \in \mathbb{F}_r} \chi(\alpha t') = 0$, then $\theta(a) = 0$. Using Proposition 3.1 we get

$$w_L(ev(a)) = 2\frac{q-1}{q}n = 2\frac{q-1}{eq}(r^2 - r). \qquad \square$$

Corollary 3.4. Let $q = 2$, the set $C_2(m, e)$ is an two-weight code with the weights 2^{2m} and $2^{2m} - 2^m$, of frequencies $2^m - 1$ and $2^{2m} - 2^m$ respectively.

Proof. If $q = 2$ then $e = 1$, by Theorem 3.3 and the sizes of M and $\mathcal{R}^*$, the proof can be easily achieved. $\square$

Theorem 3.5. Assume $\gcd(m, e) = 1$. The code $\phi(C_q(m, e))$ meets the Griesmer bound with equality.

Proof. First we recall the the Griesmer bound. Let C be an $[n, k, d]$ code over $\mathbb{F}_q$ with $k \geq 1$. Then

$$\sum_{j=0}^{k-1} \left\lceil \frac{d}{q^j} \right\rceil \leq n.$$

If $\gcd(m, e) = 1$, the code $\phi(C_q(m, e))$ have the parameters

$$n = \frac{2}{e}(q^{2m} - q^m), \quad k = 2m, \quad d = \frac{2(q-1)}{ep}(r^2 - r) = \frac{2(q-1)}{ep}(q^{2m} - q^m).$$

The ceiling function takes two values depending on j.

- $0 \leq j \leq m - 1$. In this case, $\left\lceil \frac{d}{q^j} \right\rceil = \frac{2(q-1)}{e}(q^{2m-j-1} - q^{m-j-1})$,
- $m \leq j \leq 2m - 1$. In this case, $\left\lceil \frac{d}{q^j} \right\rceil = \frac{2(q-1)q^{2m-j-1}}{e}$.

The result follows after computing two geometric series of ratio $\frac{1}{q}$. $\square$

Theorem 4.3 of [13] and Theorem 2 of [14] are just a special cases of Theorem 3.3. When $\gcd(e, m) = 1$ most the codes $\phi(C_q(m, e))$ are optimal or nearly optimal codes for given length and dimension [5]. Now we list some examples in Table IV.

Table IV: Numeral examples of Theorem 3.3

q	m	e	$[n, k, d]$	Weight distribution	Optimality
2	3	1	$[112, 6, 56]$	$1 + 56x^{56} + 7x^{64}$	Optimal
3	2	1	$[144, 4, 96]$	$1 + 72x^{96} + 8x^{108}$	Optimal
4	2	3	$[160, 4, 120]$	$1 + 240x^{120} + 15x^{128}$	Optimal
5	1	1	$[40, 2, 32]$	$1 + 20x^{32} + 4x^{40}$	Nearly Optimal
7	1	2	$[42, 2, 36]$	$1 + 42x^{36} + 6x^{42}$	Optimal
9	1	2	$[72, 2, 64]$	$1 + 72x^{64} + 8x^{72}$	Optimal

Table I: LEE WEIGHT DISTRIBUTION FOR THE CASE: $\gcd(e,m)=2$

Lee Wheight	Frequency
0	1
$2\frac{q-1}{eq}(r^2-r)$	r^2-r
$2\frac{q-1}{eq}(r^2-r^{3/2})$	$\frac{r-1}{2}$
$2\frac{q-1}{eq}(r^2+r^{3/2})$	$\frac{r-1}{2}$

3.2 The Lee weight distribution when $\gcd(e,m)=2$

Theorem 3.6. Let e be a positive divisor of $q-1$. If $\gcd(e,m)=2$ then, the set $C_q(m,e)$ is a three weight code over R with the Lee weight enumerator given in Table I.

Proof. Since $\gcd(m,e)=2$, then m is even and q is odd. If $a=0$, then $Tr(ax)=0$. So $w_L(ev(a))=0$.

If $a\in\mathcal{R}^*$ then a can be expressed as $a=\alpha+u\beta$, let $x=t+ut'\in L$. So $ax=\alpha t+u(\alpha t'+\beta t)$, and

$$Tr(ax)=Tr_{r/q}(\alpha t)+uTr_{r/q}(\alpha t'+\beta t).$$

which implies that

$$\phi(ev(a))=\phi((Tr(ax))_{x\in L})=(Tr_{r/q}(\alpha t'+\beta t),Tr_{r/q}(\alpha t'+\beta t)+Tr_{r/q}(\alpha t))_{t,t'}.$$

Taking character sum yields

$$\begin{aligned}\theta(a)&=\sum_{t\in C_0^{(e,r)}}\chi_1(Tr_{r/q}(\beta t))\sum_{t'\in\mathbb{F}_r}\chi_1(Tr_{r/q}(\alpha t'))+\sum_{t\in C_0^{(e,r)}}\chi_1(Tr_{r/q}((\alpha+\beta)t))\sum_{t'\in\mathbb{F}_r}\chi_1(Tr_{r/q}(\alpha t'))\\&=\sum_{t\in C_0^{(e,r)}}\chi(\beta t)\sum_{t'\in\mathbb{F}_r}\chi(\alpha t')+\sum_{t\in C_0^{(e,r)}}\chi((\alpha+\beta)t)\sum_{t'\in\mathbb{F}_r}\chi(\alpha t').\end{aligned}$$

Since $\sum_{t'\in\mathbb{F}_r}\chi(\alpha t')=0$, then $\theta(a)=0$. Using Proposition 3.1 we get

$$w_L(ev(a))=2\frac{q-1}{q}n=2\frac{q-1}{eq}(r^2-r).$$

If $a\in M\setminus\{0\}$, then $a=\beta u$ with $\beta\in\mathbb{F}_r^*$. By Proposition 3.2 we get

$$\sum_{s\in\mathbb{F}_q^*}\theta(sa)=4r\frac{q-1}{e}\sum_{t\in C_0^{(2,p^m)}}\chi(\beta t).$$

If $\beta\in C_i^{(2,r)}$, then we have that

$$\sum_{s\in\mathbb{F}_q^*}\theta(sa)=4r\frac{q-1}{e}\eta_i^{(2,r)}.$$

Table II: LEE WEIGHT DISTRIBUTION FOR THE CASE: $\gcd(m,e)=3$

If $p \equiv 1 \pmod 3$

Lee Wheight	Frequency
0	1
$2\frac{q-1}{eq}(r^2-r)$	r^2-r
$2\frac{q-1}{eq}(r^2-c_1r^{4/3})$	$\frac{r-1}{3}$
$2\frac{q-1}{eq}(r^2+\frac{1}{2}(c_1+9d_1)r^{4/3})$	$\frac{r-1}{3}$
$2\frac{q-1}{eq}(r^2+\frac{1}{2}(c_1-9d_1)r^{4/3})$	$\frac{r-1}{3}$

If $p \equiv 2 \pmod 3$ and $sm \equiv 2 \pmod 4$

Lee Wheight	Frequency
0	1
$2\frac{q-1}{eq}(r^2-r)$	r^2-r
$2\frac{q-1}{eq}(r^2+2r^{3/2})$	$\frac{r-1}{3}$
$2\frac{q-1}{eq}(r^2-r^{3/2})$	$2\frac{r-1}{3}$

We have that $ev(a) \in R^n$, then $\phi(ev(a)) \in \mathbb{F}_q^{2n}$, By Proposition 3.1 we get

$$4r\frac{q-1}{e}\eta_i^{(2,r)} = 2(q-1)n - qw_H(\phi(ev(a))),$$

which implies that

$$w_L(ev(a)) = 2\frac{q-1}{eq}\left(en - 2r\eta_i^{(2,r)}\right) = 2\frac{q-1}{eq}\left(r^2 - r(1+2\eta_i^{(2,r)})\right).$$

Note that $|C_0^{(2,r)}| = |C_1^{(2,r)}| = \frac{r-1}{2}$. The weight distribution of the code follows from the Lemma 2.7 and the discussion above. This completes the proof. □

Theorem 1 of [14] is a special cases of Theorem 3.6 for $e = 2$. Now we give numeral example.

Example 3.7. Let $q = 3$ and $m = 2$ and $e = 2$. Then $\phi(C_3(2,2))$ is a three-weight ternary code of parameters $[72, 4, 36]$ with the weight enumerator

$$1 + 4X^{36} + 72X^{48} + 4X^{72}.$$

3.3 The Lee weight distribution when $\gcd(e,m) = 3$

Theorem 3.8. Let e be a divisor of $q-1$. When $\gcd(m,e) = 3$ the set $C_q(m,e)$ is a code with the Lee weight distribution given in Table II, where c_1 and d_1 are uniquely given by $4q^{m/3} = c_1^2 + 27d_1^2$, $c_1 \equiv 1 \pmod 3$ and $\gcd(c_1, p) = 1$.

Proof. If $a = 0$, then $Tr(ax) = 0$. So $w_L(ev(a)) = 0$.

If $a \in \mathcal{R}^*$ then a can be expressed as $a = \alpha + u\beta$, let $x = t + ut' \in L$. a can be expressed as $a = \alpha + u\beta$. So $ax = \alpha t + u(\alpha t' + \beta t)$, and

$$Tr(ax) = Tr_{r/q}(\alpha t) + uTr_{r/q}(\alpha t' + \beta t).$$

which implies that

$$\phi(ev(a)) = \phi((Tr(ax))_{x\in L}) = (Tr_{r/q}(\alpha t' + \beta t), Tr_{r/q}(\alpha t' + \beta t) + Tr_{r/q}(\alpha t))_{t,t'}.$$

Taking character sum yields

$$\begin{aligned}\theta(a) &= \sum_{t\in C_0^{(e,r)}} \chi_1(Tr_{r/q}(\beta t)) \sum_{t'\in\mathbb{F}_r} \chi_1(Tr_{r/q}(\alpha t')) + \sum_{t\in C_0^{(e,r)}} \chi_1(Tr_{r/q}((\alpha+\beta)t)) \sum_{t'\in\mathbb{F}_r} \chi_1(Tr_{r/q}(\alpha t')) \\ &= \sum_{t\in C_0^{(e,r)}} \chi(\beta t) \sum_{t'\in\mathbb{F}_r} \chi(\alpha t') + \sum_{t\in C_0^{(e,r)}} \chi((\alpha+\beta)t) \sum_{t'\in\mathbb{F}_r} \chi(\alpha t').\end{aligned}$$

Since $\sum_{t'\in\mathbb{F}_r} \chi(\alpha t') = 0$, then $\theta(a) = 0$. Using Lemma 3.1 we get

$$w_L(ev(a)) = 2\frac{q-1}{q}n = 2\frac{q-1}{eq}(r^2 - r).$$

If $a \in M \setminus \{0\}$, then $a = \beta u$ with $\beta \in \mathbb{F}_{p^m}^*$. By the assumption $\gcd(m, e) = 3$. It then follows from Proposition 3.2 that

$$\sum_{s\in\mathbb{F}_q^*} \theta(sa) = 6r\frac{q-1}{e} \sum_{t\in C_0^{(3,r)}} \chi(\beta t).$$

If $\beta \in C_i^{(3,r)}$

$$\sum_{s\in\mathbb{F}_q^*} \theta(sa) = 6r\frac{q-1}{e}\eta_i^{(3,r)}.$$

We have that $ev(a) \in R^n$, then $\phi(ev(a)) \in \mathbb{F}_q^{2n}$, By Proposition 3.1 we get

$$6r\frac{q-1}{e}\eta_i^{(3,r)} = 2(q-1)n - qw_H(\phi(ev(a))),$$

which implies that

$$w_L(ev(a)) = 2\frac{q-1}{eq}\left(en - 3r\eta_i^{(3,r)}\right) = 2\frac{q-1}{eq}\left(r^2 - r(1 + 3\eta_i^{(3,r)})\right).$$

Since $\gcd(m, e) = 3$, then $m \equiv 0 \pmod 3$. If $p \equiv 1 \pmod 3$ by Lemma 4 the Gaussian periods $\eta_i^{(3,r)}$ take only the following three distinct values:

$$\frac{-1 + c_1 r^{1/3}}{3}, \quad \frac{-1 - \frac{1}{2}(c_1 + 9d_1)r^{1/3}}{3}, \quad \frac{-1 - \frac{1}{2}(c_1 - 9d_1)r^{1/3}}{3}$$

If $p \equiv 2 \pmod 3$ the Gaussian periods $\eta_i^{(3,r)}$ take only two distinct values. Note that $|C_0^{(3,r)}| = |C_1^{(3,r)}| = |C_2^{(3,r)}| = \frac{r-1}{3}$. The weight distribution of the code follows. This completes the proof. □

Example 3.9. Let $q = 7$ and $m = 3$ and $e = 6$. Then $\phi(C_7(3, 6))$ is a four-weight ternary code of parameters $[39102, 6, 329]$ with the weight enumerator

$$1 + 117306X^{33516} + 144X^{329} + 144X^{37044} + 144X^{30870}.$$

Table III: LEE WEIGHT DISTRIBUTION FOR THE CASE: $\gcd(e,m)=4$

If $p \equiv 1 \pmod 4$.

Lee Wheight	Frequency
0	1
$2\frac{q-1}{eq}(r^2-r)$	r^2-r
$2\frac{q-1}{eq}(r^2-r^{3/2}-2u_1r^{5/4})$	$\frac{r-1}{4}$
$2\frac{q-1}{eq}(r^2-r^{3/2}+2u_1p^{5/4})$	$\frac{r-1}{4}$
$2\frac{q-1}{eq}(r^2+r^{3/2}-4v_1r^{5/4})$	$\frac{r-1}{4}$
$2\frac{q-1}{eq}(r^2+r^{3/2}+4v_1r^{5/4})$	$\frac{r-1}{4}$

If $p \equiv 3 \pmod 4$.

Lee Wheight	Frequency
0	1
$2\frac{q-1}{eq}(r^2-r)$	r^2-r
$2\frac{q-1}{eq}(r^2-3r^{3/2})$	$\frac{r-1}{4}$
$2\frac{q-1}{eq}(r^2+r^{3/2})$	$3\frac{r-1}{4}$

Example 3.10. Let $q=4$ and $m=3$ and $e=3$. Then $\phi(C_4(3,3))$ is a three-weight code of parameters $[2688,6,1536]$ over $\mathbb{F}_4$ with the weight enumerator

$$1+12x^{1536}+4032X^{2016}+24X^{2304}.$$

3.4 The Lee weight distribution when $\gcd(e,m)=4$

Theorem 3.11. Let e be a divisor of $q-1$. Suppose that $\gcd(m,e)=4$, then the set $C_q(m,e)$ is a code with Lee weight distribution given in Table III.

Proof. Because $\gcd(m,e)=4$, then $m \equiv 0 \pmod 4$. If $p \equiv 1 \pmod 4$, then the Gaussian periods $\eta_i^{(4,p^m)}$ take only the following distinct values:

$$\frac{(1+r^{1/2}+2u_1r^{1/4})}{4}, \quad \frac{(1+r^{1/2}-2u_1r^{1/4})}{4},$$

and

$$\frac{(1-r^{1/2}+4v_1r^{1/4})}{4}, \quad \frac{(1-r^{1/2}-4v_1r^{1/4})}{4}.$$

Using a similar proof of Theorem 3.8, the proof can be easily achieved. □

Example 3.12. Let $q=5$ and $m=4$ and $e=4$. Then $\phi(C_5(4,4))$ is a four-weight code of parameters $[195000,8,142500]$ with the weight enumerator

$$1+390000X^{156000}+156X^{142500}+156X^{157500}+156X^{152500}+156X^{172500}.$$

Remark 3.13. In this paper, we describe the Lee-weight distribution of the code $C_q(m,e)$ for all e with $1 \le \gcd(e,m) \le 4$. The period polynomial $\psi_{(N,r)}(X)$ and its factorization were determined for $N=5$ by Hoshi [8], and for those dividing 8 and 12 by Gurak [6]. This means that the Lee-weight distribution of the code $C_q(m,e)$ for $\gcd(e,m) \in \{5,6,8,12\}$ can be written down now. However, the weight formulas will be messy. This demonstrates that the Lee-weight distribution of the codes $C_q(m,e)$ is indeed very complicated.

Acknowledgement. This work was supported by National Natural Science Foundation of China. Grant No.: 11171370.

References

[1] Ding, C. Linear codes from some 2-designs. IEEE Transactions on Information Theory, 2015, vol. 61, no 6, 3265-3275.

[2] Ding, K., Ding, C. Binary Linear Codes With Three Weights. IEEE Communications Letters, 2014, vol. 18, no 11, 1879-1882.

[3] Ding, C., Li, C., Li, N., Zhou, Z. Three-weight cyclic codes and their weight distributions. Discrete Mathematics, 2016, vol. 339, no 2, 415-427.

[4] Ding, C., Luo, J., Niederreiter, H. Two-weight codes punctured from irreducible cyclic codes. In : Proc. of the First International Workshop on Coding Theory and Cryptography. 2008. 119-124.

[5] Grassl, M. Bounds on the minimum distance of linear codes and quantum codes. Online available at http://www.codetables.de.

[6] Gurak, S. J. Period polynomials for $\mathbb{F}_q$ of fixed small degree. CRM Proc. and Lect. Notes, 2004, vol. 36, 127-145.

[7] Heng, Z., Yue, Q. A class of binary linear codes with at most three weights. IEEE Communications Letters, 2015, vol. 19, no 9, 1488-1491.

[8] Hoshi, A. Explicit lifts of quintic Jacobi sums and period polynomials for $\mathbb{F}_q$. Proceedings of the Japan Academy, Series A, Mathematical Sciences, 2006, vol. 82, no 7, 87-92.

[9] Lidl, R., Niederreiter, H. Finite fields. Cambridge university press, 1997.

[10] Ma, C., Zeng, L., Liu, Y., Feng, D., Ding, C. The weight enumerator of a class of cyclic codes. IEEE Transactions on Information Theory, 2011, vol. 57, no 1, 397-402.

[11] Myerson, G. Period polynomials and Gauss sums for finite fields. Acta Arithmetica, 1981, vol. 39, no 3, 251-264.

[12] Shi, M. J., Liu, Y., Solé, P. Optimal two weight codes from trace codes over a non-chain ring. Discrete Applied Mathematics,(Accepted), 2016.

[13] Shi, M., Luo, Y., Solé, P. Optimal Two-Weight Codes From Trace Codes Over $\mathbb{F}_2 + u\mathbb{F}_2$. IEEE Communications Letters, 2016, vol. 20, no 12, 2346-2349.

[14] Shi, M., Luo, Y., Solé, P. Construction of one-Lee weight and two-Lee weight codes over $\mathbb{F}_p + v\mathbb{F}_p$. Journal of Systems Science and Complexity, 2016, 1-10.

[15] Yu, L., Liu, H. The weight distribution of a family of p-ary cyclic codes. Designs, Codes and Cryptography, 2016, vol. 78, no 3, 731-745.